

# **Evolution of Hematopoietic Stem Cell Transplant Programs in Malaysia**

Norashikin Saidon<sup>1</sup>, Nur Adila Anuar<sup>2</sup>, Chang Kian Meng<sup>3</sup>, Ong Tee Chuan<sup>4</sup>, Tan Sen Mui<sup>4</sup>, Gan Gin Gin<sup>2</sup>, Fadilah SAW<sup>5</sup>, Hany Ariffin<sup>6</sup>, Lin Hai Peng<sup>7</sup>, Chan Lee Lee<sup>7</sup>, Alan Teh<sup>7</sup>, Bee Ping Chong<sup>2</sup>

<sup>1</sup>Faculty of Medicine, Sungai Buloh, Universiti Teknologi MARA, Malaysia, <sup>2</sup>Hematology Unit, University of Malaya Medical Center, Malaysia, <sup>3</sup>Sunway Medical Center, Malaysia, <sup>4</sup>Hematology Department, Hospital Ampang, Malaysia, <sup>5</sup>Department of Medicine, Universiti Kebangsaan Malaysia Medical Centre, Malaysia, <sup>6</sup>Department of Pediatrics, University of Malaya Medical Center, Malaysia, <sup>7</sup>Subang Jaya Medical Center, Malaysia

### Abstract

Hematopoietic stem cell transplantation (HSCT) is now widely practiced worldwide. It has the potential to cure many hematological diseases, such as acute leukemia and thalassemia. As an emerging country, Malaysia has made advancements despite many challenges. HSCT has evolved rapidly since the first pediatric allogeneic HSCT case in 1987. The first adult HSCT was performed 5 years later in 1993. Currently, a total of 13 hospitals offer HSCT services throughout Malaysia. These include private healthcare services, substantially funded government hospitals governed by the Ministry of Health, and partially funded teaching hospitals governed by the Ministry of Education. Until 2015, 1,987 allogeneic and 1,648 autologous HSCT procedures were performed. This article narrates the history and development of HSCT in Malaysia and briefly discusses the challenging issues in this area.

Key words: hematopoietic stem cell transplant, Malaysia, allogeneic, autologous, haploidentical

Submitted December 3, 2019; Accepted March 6, 2020; Published online June 10, 2020; Issued online August 25, 2020 Correspondence: Bee Ping Chong, Haemotology Unit, Department of Medicine, Faculty of Medicine, University Malaya Kuala Lumpur 50603, Malaysia, E-mail: bpingchong@gmail.com

# Introduction

The first pediatric hematopoietic stem cell transplantation (HSCT) was performed at the University of Malaya Medical Center in 1987, followed by an adult HSCT in 1993<sup>1,2</sup>. Since then, HSCT in Malaysia has rapidly evolved. At present, a total of 13 hospitals offer HSCT services throughout Malaysia, including private healthcare services, substantially funded government hospitals governed by the Ministry of Health, and partially funded teaching hospitals governed by the Ministry of Education. Twelve of the centers are within the Malaysian Peninsular and the other is in the Malaysian Borneo (**Table 1**).

# Malaysian National Donor Program

The Malaysian Stem Cell Registry (MSCR) was first established in 2000. It was a collaborative effort between the Ministry of Health (Medical Development Division and Institute of Medical Research) and the National Cancer Council Malaysia (MAKNA)<sup>1</sup>. The registry aims to promote and recruit donors for bone marrow and peripheral stem cell transplantation. To date, 31,200 donors have been registered. In 2003, the Cord Blood Bank Unit was established alongside the MSCR. This was based on convincing evidence for the use of hematopoietic stem cells, sourced from umbilical cord blood, as an alternative to bone marrow transplantation, especially for hematological, immunological, and metabolic storage disorders in children and young adults. Cord blood units (CBU) help overcome shortages for patients requiring HSCT, especially those having no suitable family members or having unrelated donors<sup>2</sup>. CBUs are obtained, with consent, from maternity wards in four hospitals within Kuala Lumpur and a single hospital in Alor Setar. There are 7,450 CBUs available at present. In 2014, the National Stem Cell Coordinating Center, which is based at the National Blood Bank, was established to better facilitate HSCT from unrelated donors. It provides a one-stop center for identification and coordination of transplantation from donors and CBUs and facilitates the matching process among unrelated stem cell transplant donors and receivers.

YEAR	20	15	1987-2015		
NAME OF HOSPITALS	n	%	n	%	
HOSPITAL AMPANG		60	1,466	40	
HOSPITAL KUALA LUMPUR, INSTITUTE PAEDIATRICS	27 7		449	12	
HOSPITAL UNIVERSITI KEBANGSAAN MALAYSIA	14	3	246	7	
SUBANG JAYA MEDICAL CENTRE (ADULT)	37	9	412	11	
SUBANG JAYA MEDICAL CENTRE (PAED)	20	5	106	3	
UNIVERSITY OF MALAYA MEDICAL CENTRE (ADULT)	16	4	273	8	
UNIVERSITY OF MALAYA MEDICAL CENTRE (PAED)	15	4	446	12	
GLENEAGLES MEDICAL CENTRE, PENANG	0	0	21	1	
LAM WAH EE HOSPITAL, PENANG	0	0	9	0	
HOSPITAL UNIVERSITI SAINS MALAYSIA	0	0	46	1	
AMPANG PUTERI SPECIALIST HOSPITAL	0	0	6	0	
HOSPITAL PULAU PINANG	28	7	138	4	
SABAH WOMEN'S AND CHILDREN'S HOSPITAL	0	0	2	0	
HOSPITAL MELAKA	3	1	3	0	
TOTAL	401	100	3,626	100	

Table 1. List of hospitals with haematopoietic stem cell transplant services, 1987-2015



Figure 1. New transplant rate per million population (pmp), 1987-2015

# Indications and Characteristics of HSCT in Malaysia

The number of HSCTs in Malaysia has increased rapidly over the years (**Figure 1**)<sup>3</sup>. In 2015, 401 HSCTs were performed (**Table 2**)<sup>3</sup>. The majority of HSCTs involved patients diagnosed with acute leukemia and lymphomas, followed by plasma cell dyscrasia and hemoglobinopathies. A total of 3,626 HSCTs were performed from 1987 through 2015; 1,978 allogeneic and 1,648 autologous HSCTs. The majority (78%) of hematopoietic stem cells (HSC) were sourced from a combination of marrow stem cells with peripheral blood or peripheral blood alone. Other sources of HSC include the bone marrow (19%) and both bone marrow with cord blood (3%) or cord blood alone<sup>3,4</sup>.

The majority of HSCTs involve Human Leukocyte Antigen (HLA)-matched siblings (88%), followed by matched unrelated donors (3%). The remainder includes syngeneic transplantation and related and unrelated HLA-mismatch donors. In 2016, 148 allogeneic and 212 autologous HSCTs were performed. Notably, this was the year that haploidentical family donors surpassed matched unrelated donors. The use of haploidentical family donors as an alternative source of HSCs in Malaysia has followed the trend observed in other countries i. e., an increase in the number of cases. Haploidentical HSCT has enabled all patients to find a donor and compared to matched unrelated donors (MUD), it also has the advantages of lower cost and faster graft acquisition. In addition, outcomes from haploidentical HSCTs have also greatly improved over the years with better methods preventing graft versus host disease and the increasing experience of transplant center personnel.

The survival of HSCT patients has improved over the years (**Figure 2**)<sup>3</sup>. This is because of the availability of more effective drugs and more accurate diagnostic tests. Compared to the  $\leq 60\%$  survival rate of HSCT patients in the 1990s, the survival rate of HSCT patients was approximately 70% between 2010 and 2015<sup>3</sup>. The main cause of death between 1987 and 2015 was the underlying disease itself, followed by sepsis, graft versus host disease, hemorrhage, venous occlusive disease, and organ failure.

#### **HSCT** Issues in Malaysia

There are multiple issues pertaining to HSCT in Malaysia, especially funding for the transplant. Other issues include availability of both laboratory facilities and unrelated donors. The numbers of voluntary donors in our local registry remains small. More efforts are needed to promote awareness and to encourage volunteer donors. Education and proper counseling at the time of prerecruitment remains pertinent to ensure donors are fully

Table 2. Indications for Haematopoietic stem cell transplant from 2010-2015

YEAR	2010	2011	2012	2013	2014	2015	TOTAL
DIAGNOSIS	n	n	n	n	n	n	n
Acute Leukemias	105	84	87	99	104	117	596
Chronic Myelogenous Leukemia (CML)	15	15	10	6	23	25	94
Other leukemias	1	1	0	2	0	1	5
Lymphomas	65	99	115	127	124	154	684
Myelodysplastic Syndrome (MDS)	4	5	8	4	9	9	39
Combined Myelodysplastic/Myeloproliferative Syndrome (MD/MPS)	0	0	2	0	0	0	2
Myeloproliferative Syndrome	1	1	0	0	3	6	11
Plasma cell Disorder including Multiple Myeloma	23	29	47	45	45	47	236
Anaemia	14	20	10	13	10	15	82
Hemoglobinopathy	13	10	8	12	14	11	68
Other Malignancies	16	5	3	6	8	13	51
Primary Immune Deficiencies	1	1	0	0	2	0	4
Inherited Disorders of Metabolism	1	0	0	2	1	2	6
Platelet and Other Inherited Disorders	1	0	0	0	0	0	1
Histiocytic Disorders	0	1	13	1	0	1	16
TOTAL	260	271	303	317	343	401	1,895



Figure 2. Patient survival by year of transplant 1987-2015

committed. Procurement of stem cells from MUD is not financially supported by our government at this currently, although the transplantation itself is heavily subsidized in government hospitals. Increasingly, patients and hospitals are opting for haploidentical related donors as an alternative because of the high cost of procurement of MUD stem cells from overseas.

Appropriate laboratory facilities are essential for HSCT. A pre-transplant workup includes HLA typing for both patient and donor. High resolution HLA typing was previously only available at the Institute of Medical Research. Recently in 2018, the University of Malaya Medical Center begun offering this service in collaboration with an overseas laboratory. Additionally, as posttransplant viral infection remains a common complication, identifying and monitoring such infection is essential. Unfortunately, the facilities and expertise required for this remain limited to a few centers. Financial constraints have affected the training and education of service providers as well as the setting up of necessary facilities. A unique, albeit complex, trait of our healthcare system includes the provision of service by three major sectors: the Ministry of Health, Ministry of Education (MOE), and private healthcare services. Thus, services surrounding stem cell transplantation are unsynchronized, as subsidization and source of financial assistance differ between these sectors. For example, under the MOE, patients requiring stem cell transplantation may only be partially subsidized if deemed eligible. Generally, the cost of transplantation is estimated at RM 50,000 or USD 15,000. Further financial assistance, if required, often depends on eligibility for social welfare support. Conversely, the cost in the private sector is much higher, up to RM 200,000 or USD 50,000.

#### Conclusion

Over time, HSCT outcomes have markedly improved and are presently very encouraging. With the increase in transplant centers throughout the country, the number of transplantations in Malaysia is expected to rise in the future. Haploidentical donor transplantations that have begun in a few centers will further rapidly increase in number. The future of transplantation in Malaysia could be further improved if the stakeholders would commit to resolving the issues, especially those of a financial nature, which we are currently facing.

#### Acknowledgments

We thank Dr. Chitra from National Stem Cell Coordinating Center and Dr. Naren Kumar from National Transplant Registry for providing the data.

#### Author's Contribution

N. S and N. A. A wrote the manuscript with the support and supervision from B. P. C and G. G. G. B. P. C, G. G. G. C. K. M, O. T. C, T. S. M, F. SAW, H. A, L. H. P, C. L. L and A. T contributed to the information and data to write the manuscript.

#### Conflict of Interest

The authors declare no conflict of interest. Disclosure forms provided by the authors are available here.

#### References

- Gan GG, Teh A, Chan LL, Cheong SK, Chang KM, Ibrahim H. Bone marrow and stem cell transplantation: Malaysian experience. Bone Marrow Transplant. 2008; 42 Suppl 1: S103-S5.
- Fadilah SAW, Leong CF, Cheong SK. Stem cell transplantation in Malaysia and future directions. Med J Malaysia. 2008; 63: 279-80.
- Hooi LS, Chandramalar S, editors. Twelfth Report of the National Transplant Registry 2015. Kuala Lumpur: National Transplant Registry; 2018.
- Ernest M, Nyi NN, Bachok N, Husin A. Survival and prognostic factors in Malaysian acute myeloid leukemia patients after allogeneic haemtopoietic setm cell transplantation. Int J Hematol. 2013; 98: 197-205.

https://doi.org/10.31547/bct-2019-017 Copyright © 2020 APBMT. All Rights Reserved.